

**FEEDING HABITS OF THE BIGEYE SCAD,
SELAR CRUMENOPHTHALMUS (CARANGIDAE),
IN LA RÉUNION ISLAND WATERS
(SOUTH-WESTERN INDIAN OCEAN)**

by

Olivier ROUX (1) & François CONAND (2)

ABSTRACT. *Selar crumenophthalmus*, the bigeye scad, is a coastal pelagic fish, which feeds mostly at night on large planktonic animals. Young scads prey mostly on crustaceans (euphausiids, decapods) and adults are active predators of fish larvae and early juveniles. The large size of the eyes could be an adaptation to night predation.

RÉSUMÉ. Comportement alimentaire du pêche-cavale, *Selar crumenophthalmus* (Carangidés) sur les côtes de La Réunion (sud-ouest de l'océan Indien).

Selar crumenophthalmus est un poisson pélagique côtier prédateur qui se nourrit, surtout la nuit, de la fraction la plus grosse du plancton. Les jeunes individus consomment principalement des crustacés (Euphausiacés, Décapodes) et les adultes sont surtout prédateurs de larves et juvéniles de poissons. La grande taille des yeux pourrait être une adaptation à la chasse nocturne.

Keywords. Carangidae - *Selar crumenophthalmus* - SW - La Réunion I. - Feeding - Predation - Diet.

The bigeye scad, *Selar crumenophthalmus* (Bloch, 1793), is a small coastal pelagic fish, which is common in the tropical and subtropical belt of all oceans. This species is fished either for human consumption or as bait for tuna fishing. In some places it can be of major importance to fisheries and seems to be especially important in the coastal waters of islands with no continental shelf and no large populations of clupeids. This is the case in La Réunion Island where it is the most important small pelagic fish, with the annual catch estimated at 100 tonnes.

The biology of the bigeye scad has been studied in many different countries, the most recent of which were conducted in Hawaii (Clarke and Privitera, 1995; Iwai *et al.*, 1996) and the Philippines (Dalzell and Penaflo, 1989). Some other studies on coastal pelagic fisheries include this species, but little research has been carried out on the ecology of the bigeye scad. In 1994, Roux studied the growth and biology of the bigeye scad in La Réunion Island, showing that this species has an annual reproductive cycle. Spawning occurs mostly from October to December with the length of one-year old individuals averaging 215 mm (fork length). Few survive longer than one year.

The importance of the bigeye scad in the food chain of coastal areas and the prospect of possible culture development prompted this detailed first study into the nutrition and feeding behaviour of that species.

(1) FAO, FIDI, viale delle Terme di Caracalla, 00100 Roma, ITALIA. [olivier.roux@fao.org]

(2) Université de La Réunion, ECOMAR, 97715 St Denis cedex 9, LA RÉUNION.
[fconand@univ-reunion.fr]




Fig. 1. Location of La Réunion Island and of sampling stations.

MATERIAL AND METHODS

Sampling and prey analyses

Bigeye scads were sampled from fishermen catches in two locations where the fishing gear is different (Fig. 1). At night, in the Bay or in the harbour of La Possession, fish are attracted by light and caught with lines from a skiff. In Saint Paul Bay, scads are caught during the day, usually in the morning, with a beach seine. Although sampling of fish caught with a beach seine is relatively simple, the fish usually remain at least one hour in the net during the fishing operation during which time regurgitation may occur, biasing the observations. Therefore, only samples resulting from very fast operations were used for this study.

The content of a total of 182 stomachs was examined. For hand line night fishing, between 4 and 5 a.m., a total of 138 scads were collected, from February to May. For beach seine during the light, between noon and 2 p.m., 44 scads were collected in 4 samples at various months of the year.

Fork length was measured, and stomachs - from oesophagus to pylorus - were collected from freshly killed fish. They were preserved in 8% formalin. The number of stomachs of each sample varied between 15 and 20. This sample size is recommended by Bell and Harmelin-Vivien (1983) who showed that beyond this number no new prey category usually appears and the diversity index reaches a platform. The different prey categories were sorted by taxonomic groups for each stomach contents and further determination was made whenever possible.

For prey counting, each individual was considered as an item, except for colonial organisms or algae, which were counted as one item per colony. Unidentifiable prey fragments found in stomachs were also considered as one item. Each prey category for each stomach was weighed separately, at a precision level of 0.1 mg. This wet weight was measured after wiping the prey on soft filter paper, as recommended by Windell (1971). Using that wet weight allows for a better account of the real bulkiness of prey items in the digestive tract.

Feeding activity

Various methods can be used to estimate the feeding activity. Basic information is given by the percentage of empty stomachs, E_r :

$$E_r = \frac{\text{number of empty stomachs}}{\text{number of stomachs sampled}}$$

Other useful information is the repletion level of the stomach, given by the repletion index, R_i :

$$R_i = \frac{\text{weight of stomach contents (g)}}{\text{fork length of fish (mm)}}$$

This index was originally defined by Berg (1979), who used the weight of the fish instead of its fork length. The use of length avoids the influence of the fat level, the state of the gonads and the weight of the food in the stomach.

Determination of diet

Three primary indices N , W and P were calculated for each prey item (I) and for each size class of fish:

$$N_i = \frac{\text{number of individuals of item } I}{\text{total number of prey}}$$

$$W_i = \frac{\text{weight of one item } I}{\text{total weight of prey}}$$

$$F_i = \frac{\text{number of stomachs containing item } I}{\text{number of non empty stomachs examined}}$$

Following Zander (1982), the Main Food Item index (MFI), which combines these 3 indices, was used for each item:

$$MFI_I = \sqrt{\frac{N_i + F_i}{2}} \times W_i$$

The use of this index was improved by Rosecchi and Nouaze (1987) in the following way. The MFI was summed and for each prey item was expressed as the ratio (%) of the total. Prey items are then classified by decreasing relative values and cumulated ratios are calculated. From these cumulated ratios, they proposed the following classification:

main prey:	the prey of the first rank, and cumulated ratio MFI $\geq 50\%$
secondary prey:	cumulated ratio MFI between 50 and 75%
accessory prey:	cumulated ratio MFI $\geq 75\%$

RESULTS

Feeding activity

Fish caught during the day often have empty stomachs with a mean empty ratio of $E_{r(\text{day})} = 38.32\%$, confidence interval at $\alpha = 0.05$. At night most of the fish have food in their stomachs and the mean empty ratio is $E_{r(\text{night})} = 11.38\%$. Considering the non empty stomachs only, the mean repletion index with confidence interval at $\alpha = 0.05$, is $R_{i(\text{day})} = 0.0232\%$ during the day, and $R_{i(\text{night})} = 0.4930\%$ at night. These two results show that most individuals are feeding actively at night.

Diet

The food found in the stomachs was only composed of pelagic animals. No mineral material (sand, shell fragments), algae or benthic animals were found. Ten prey categories were observed (Table II). The analysis shows that the diet composition varies with scad size. Hence MFI was computed by size classes with 20 mm intervals. Preliminary results showed that some size classes have similar diets and could be pooled. Three size classes were retained: 100-119 mm, 120-159 mm, and 160-219 mm. The contribution of each category, expressed as percentages of total MFI and cumulated MFI, are given in table III. It appears that: (1) small scads feed mainly on euphausiids, shrimps and crabs, (2) crabs and fish are the main prey of medium size fish, and (3) fish are the main prey of the larger individuals (Table III). Shrimps and other crustaceans are the secondary categories of prey in all size classes.

The fish found in stomach contents were mostly juveniles of clupeids and pelagic post larvae of balistids and serranids. Their size was usually between 1 and 5 cm. The crabs were megalops larvae of various unidentified species. All the other food items were also pelagic animals from the macroplankton.

DISCUSSION

The study of feeding activity of *Selar crumenophthalmus* in La Réunion Island, shows that empty stomachs are common during the day but not at night, indicating that the bigeye scad mainly feeds during the night. The same observation was made in Hawaii by Kawamoto (1973) and by Tobias (1987) in the Virgin Islands. Pillay (1952) stressed the fact that the stomach repletion level can be biased by the sampling method because fish can regurgitate, or ingest food during capture. Given that night samplings using line and hooks do not show regurgitation, the observation of night feeding cannot be refuted. For fish caught with the beach seine, there is no proof that there is no regurgitation and the assumption of little feeding during the day has still to be investigated.

Table II. Prey items found in the stomachs of *Selar crumenophthalmus* in La Réunion Island.

Prey items	Mean weight (g) with confidence interval at $\alpha = 0.05$	Remarks
Fish	0.457 ± 0.090	Mostly, larvae and juveniles of fish: Clupeidae, Balistidae, Serranidae, Leptocephals.
Crab larvae	0.041 ± 0.009	Planktonic megalops.
Euphausiids	0.009 ± 0.002	
Shrimps	0.070 ± 0.024	Swimming species of decapods.
Stomatopods	0.214 ± 0.026	Planktonic larvae of Squillidae.
Amphipods	0.004 ± 0.001	Planktonic animal living in patches; only 4 occurrences, but great number of individuals.
Crustacean fragments	0.101 ± 0.021	Mainly chitinous debris.
Coleoptera	0.018 ± 0.001	From terrestrial origin.
Cephalopods	0.293 ± 0.061	Mostly octopus.
Unidentified	0.132 ± 0.059	

Night feeding is not common for epipelagic fish although it is not an exception. Small plankton feeders, like sprats and anchovies, can feed at night (Milton *et al.*, 1990). Predators like tunas and large pelagic fish, usually do not prey at night (Reintjes and

Table III.1. Prey classification obtained with the Main Food Item index (MFI) in the three defined size classes of *Selar crumenophthalmus* in La Réunion Island.

100-119 mm (n = 8)	MFI	% of total MFI	% of cumulated MFI
Euphausiids	0.715	59.6	59.6
Shrimps	0.152	12.7	72.3
Crab larvae	0.119	9.9	82.1
Stomatopods	0.095	7.9	90.1
Crustacean fragments	0.071	5.9	96.0
Unidentified	0.049	4.0	100
Fish	0	0	100
Amphipods	0	0	100
Coleoptera	0	0	100
Cephalopods	0	0	100
120-159 mm (n = 59)	MFI	% of total MFI	% of cumulated MFI
Crab larvae	0.345	28.8	28.8
Fish	0.214	17.9	46.7
Shrimps	0.176	14.7	61.4
Euphausiids	0.155	13.0	74.4
Crustacean fragments	0.110	9.2	83.6
Stomatopods	0.104	8.7	92.3
Cephalopods	0.046	3.5	95.8
Unidentified	0.042	3.8	99.6
Amphipods	0.002	0.2	99.8
Coleoptera	0.002	0.2	100
160-219 mm (n = 63)	MFI	% of total MFI	% of cumulated MFI
Fish	0.464	40.1	40.1
Crab larvae	0.200	17.2	57.3
Shrimps	0.140	12.1	69.3
Crustacean fragments	0.120	10.4	79.7
Cephalopods	0.076	6.6	86.3
Stomatopods	0.062	5.4	91.7
Unidentified	0.052	4.4	96.1
Amphipods	0.036	3.1	99.3
Euphausiids	0.007	0.6	99.9
Coleoptera	0.002	0.1	100

Table III.1.2. Main and secondary prey in the three size classes of the *Selar crumenophthalmus* in La Réunion Island.

Size classes	Main prey categories	Secondary prey categories
100 - 119 mm	Euphausiids	Shrimps
120 - 159 mm	Crab, fish	Shrimps, euphausiids
160 - 219 mm	Fish	Crab, shrimps

King, 1953; Buckley and Miller, 1994), because these fish need to see their prey for the chase (Marsac and Cayré, 1998). Exceptions however were observed when there was a bright moon light, which confirms the need of light for predation in tunas. The comparatively large size of the eyes of the bigeye scad probably explains their ability to prey at night in the dark and particularly on slow swimming animals from the macroplankton.

The composition of the food of *Selar crumenophthalmus* showed that this species is zooplanktivorous as a juvenile and predominantly piscivorous when adult. The absence of mineral material excludes the possibility of a benthic food intake (Harmelin-Vivien, 1979). In some instances, the occurrence of insects in the diet confirms the general observation of fishermen that *S. crumenophthalmus* often feeds near the surface. This predatory behaviour is characteristic of small carangids and was also observed in *Trachurus* spp. (Ben Salem, 1988).

The catch of small coastal pelagic fish in La Réunion Island waters is estimated to 160 tonnes (catch statistic for 1997 from the fishery administration). The bigeye scad is the most common fish with a catch of 100 tonnes, other scads contributing with 50 tonnes, and sardines representing only 10 tonnes. The absence of a continental shelf, estuaries and protected areas like other volcanic islands (Cabo Verde, Hawaii), could be an explanation of the low abundance of sardines, which feed mainly on phytoplankton and small zooplankton. The opposite is generally observed around islands with an extensive continental shelf.

Acknowledgements. This study was financially supported by the University of La Réunion, IFREMER and IRD. The authors wish to thank D. Ross for his advices and help during the field study, E. Tessier for the information he provided on the fishery, and J.R. Enilorac master fisherman and his team.

REFERENCES

- BELL J.D. & M.L. HARMELIN-VIVIEN, 1983. Fish fauna of French Mediterranean *Posidonia oceanica* seagrass meadows. Feeding habits. *Tethys*, 11(1): 1-14.
- BEN SALEM M., 1988. Régime alimentaire de *Trachurus trachurus* (Linnaeus, 1758) et de *T. mediterraneus* (Steindachner, 1868), (Poissons, Téléostéens, Carangidés) de la province atlantico-méditerranéenne. *Cybius*, 12(3): 247-253.
- BERG J., 1979. Discussion of methods of investigating the food of fishes, with reference to a preliminary study of the food of *Gobiusculus flavescens*. *Mar. Biol.*, 50: 263-273.
- BUCKLEY T.W. & B.S. MILLER, 1994. Feeding habits of yellowfin tuna associated with fish aggregation devices in American Samoa. *Bull. Mar. Sci.*, 55(2-3): 445-459.
- CLARKE T.A. & L.A. PRIVITERA, 1995. Reproductive biology of two Hawaiian pelagic carangid fishes, the bigeye scad, *Selar crumenophthalmus*, and the round scad, *Decapterus macarellus*. *Bull. Mar. Sci.*, 56(1): 33-47.
- DALZELL P. & G. PENAFLORE, 1989. The fisheries biology of the bigeye scad *Selar crumenophthalmus* in the Philippines. *Asian Fish. Sci.*, 3: 115-131.
- HARMELIN-VIVIEN M.L., 1979. Ichtyofaune des récifs coralliens de Tuléar (Madagascar): écologie et relations trophiques. Thèse de Doctorat d'État, 165p. Univ. Aix-Marseille II.
- IWAI T.Y. Jr., TAMARU C.S., YASUKOCHI L., ALEXANDER S., YOSHIMURA R. & M. MITSUYATSU, 1996. Natural spawning of captive bigeye scad *Selar crumenophthalmus* in Hawaii. *J. World Aquacult. Soc.*, 27(3): 332-339.
- KAWAMOTO P.Y., 1973. Management investigation of the akule or bigeye scad, *Trachurus crumenophthalmus* (Bloch). Hawaii Division of Fisheries and Game, Project Report N° H-4-R: 28p.

- MARSAC F. & P. CAYRÉ, 1998. **T**elemetry applied to behaviour analysis of yellowfin tuna (*Thunnus albacares*, Bonnaterre, 1788) movements in a network of fish aggregating devices. *Hydrobiologia*, 371/372: 155-171.
- MILTON D.A., BLABER S.J.M. & N.J.F. RAWLINSON, 1990. **D**iet and prey selection of six species of tuna baitfish in three coral reef lagoons in the Solomon Islands. *J. Fish Biol.*, 37: 205-224.
- PILLAY T.V.R., 1952. **A** critic of the methods of study of food of fishes. *J. Zool. Soc. India*, 4: 185-200.
- REINTJES J.W. & J.E. KING, 1953. **F**ood of yellowfin tuna in the Central Pacific. *Fish. Bull. U.S.*, 54(81): 91-110.
- ROSECCHI E. & Y. NOUAZE, 1985. **C**omparaison de cinq indices alimentaires utilisés dans l'analyse des contenus stomacaux. *Rev. Trav. Inst. Pêches marit.*, 49(3-4): 111-123.
- ROUX O., 1994. **B**iology and ecology of the cavale (*Selar crumenophthalmus*) at the island of La Réunion. Mémoire stage DEA, 40p. Univ. Aix-Marseille.
- TOBIAS W., 1987. **R**esource evaluation of the bigeye scad *Selar crumenophthalmus* in the insular shelf waters around St Croix, US Virgin Islands. *Proc. Gulf. Carib. Fish. Inst.*, 40: 82-98.
- WINDELL J.T., 1971. **F**ood analysis and rate of digestion. In: *Methods for Assessment of Fish Production in Freshwaters* (Ricker W.E., ed), pp. 15-226. Oxford: Blackwell.
- ZANDER C.D., 1982. **F**eeding ecology of littoral gobiid and blennioid fish of the Banyuls area (Mediterranean Sea). I. Main food and trophic dimension of niche and ecotope. *Vie Milieu*, 32: 1-10.

Reçu le 06.12.1999.

Accepté pour publication le 20.04.2000.